

# Body Height and its Estimation Utilizing Arm Span Measurements in Male Adolescents from Southern Region in Montenegro

**Rajko Milašinović and Stevo Popović**

University of Montenegro, Faculty for Sport and Physical Education, Nikšić, Montenegro

**Radenko Matić**

University of Novi Sad, Faculty of Sport and Physical Education, Novi Sad, Serbia

**Jovan Gardašević and Duško Bjelica**

University of Montenegro, Faculty for Sport and Physical Education, Nikšić, Montenegro

## ABSTRACT

*The purpose of this study was to examine the body height in Montenegrin male adolescents from southern region as well as the relationship between arm span as an alternative to estimating the body height, which would vary from region to region in Montenegro. Our investigation analyses 87 male adolescents from the southern region in Montenegro. The anthropometric measurements were taken according to the protocol of the International Society for the Advancement of Kinanthropometry (ISAK). Means and standard deviations regarding the anthropometric measurements were obtained. The relationships between body height and arm span were determined using simple correlation coefficients and their 95% confidence interval. Then a linear regression analysis was performed to examine the extent to which the arm span can reliably predict body height. The results displayed that male Southern-Montenegrins are  $182.53 \pm 7.53$  cm tall and have an arm span of  $184.55 \pm 9.03$  cm. Compared to other studies, the results of this study have shown that this gender made Southern-Montenegrins the tall population, taller than most of nation around the Europe. On the other hand, expectably, the arm span reliably predicts body height in this gender. However, the estimation equations which have been obtained in Southern-Montenegrins are, different alike in general population, since arm span was closer to body heights ( $2.03 \pm 1.50$  cm), more than in general population. Hence, this study also confirms the necessity for developing separate height models for each region in Montenegro.*

**Key words:** stature, armspan, region, boys, Montenegro

## Introduction

The unusual tallness of Montenegrin highlanders was a fact recognized by European anthropologists more than 100 years ago (Bjelica, Popović, Kezunović, Petković, Jurak, & Grasgruber, 2012). A sample of 800 Montenegrin men measured by Robert W. Ehrich (Coon, 1975) at the beginning of the 20th century gave the highest average in all of Europe (177 cm), with some districts approaching 178 centimeters. Furthermore, a contemporary research study conducted by Pineau, Delamarque, and Božinović (2005), showed that the male population of the Dinaric Alps is on average, the tallest in whole Europe. Thus, this study has challenged many scientists to believe that Montenegrins are still the tallest population in Europe and Bjelica and his collaborators (2012) confirm they are very tall but not the tallest with 183.2 centimeters. From the reason the sample in this study was created by university students, a more recent study was conducted as a national survey (Popović, Bjelica, & Hadžić, 2014) and it confirmed the results of the previous study and found the average body height of Montenegrin male adolescents were 183.74 centimeters tall. From this reason the modern Montenegrins fall partly into the Dinaric racial classification (Bjelica et al., 2012), the authors did believe the male population that live in the southern region might be much taller than average Dinaric Alps subjects, mostly due to the much

better lifestyle in the coastal area. Hence, the purpose of this study was twofold. The first purpose was to examine the body height in Montenegrin male adolescents from southern region as the authors did believe this is the place where the population may reach the full potential of the Dinaric Alps, while the second purpose was to examine the relationship between body height and arm span as an alternative to estimating the body height, which would vary from region to region in Montenegro.

## Methods

The nature and scope of this study qualifies 87 male adolescents from the southern region in Montenegro to be subjects. The average age of the male subject was  $18.29 \pm 0.63$  years old (range 17-19 yrs). It is also important to emphasize that the authors could not accept adolescents with physical deformities that could affect body height or arm span, and without informed consent were excluded from the study. The exclusion criterion was also being non-Southern Montenegrin.

According to Marfell-Jones, Olds, Stew, and Carter (2006), the anthropometric measurements, including body height and arm span were taken according to the protocol of the International Society for the Advancement of Kinanthropometry (ISAK). The trained anthropometrist (the same one for each measure)

whose quality of performance was evaluated against prescribed “ISAK Manual” prior to the study performed these measurements. The age of the individuals was determined directly from their reported date of birth.

The body height presents the perpendicular distance between the top of the head (the vertex) and the bottom of the feet. It was measured using stadiometer to the nearest 0.1 centimeters in bare feet with the participants standing upright against a stadiometer. The respondents had to put their feet together and move back until their heels touched the bottom of the stadiometer upright. Their buttocks and upper part of their back have also been touching the stadiometer upright while their head didn’t have to touch the stadiometer. The respondent’s head had to be in the Frankfort horizontal plane. This was achieved when the lower edge of the eye socket (the orbitale) is horizontal with the tragion. The vertex was the highest point on their head, otherwise the respondents had to raise or lower their chin until it was in the Frankfort horizontal plane to align their head properly.

The arm span is the anthropometric measurement of the length from the tip of the middle fingers of the left and right hands when raised parallel to the ground at shoulder height at a one-hundred eighty degree angle. It was measured using a calibrated steel tape to the nearest 0.1 centimeters in bare feet on a level concrete floor with their upper backs, buttocks and heels against the wall which provide support. The participant’s head was also in the Frankfort horizontal plane and the arms were outstretched at right angles to the body with palms facing forwards. The measurement were taken from one middle fingertip

to the other middle fingertip, with the tape passing in front of the clavicles while two field workers supported the elbows. The measurements were taken twice, and an average of the two readings was calculated. When the two measurements agreed within 0.4 centimeters, their average was taken as the best estimate for the true value. When the two initial measures didn’t satisfy the 0.4 centimeters criterion, two additional determinations were made and the mean of the closest records was used as the best score.

The analysis was carried out using Statistical Package for Social Sciences (SPSS) version 20.0. Means and standard deviations (SD) were obtained for both anthropometric variables. A comparison of means of body heights and arm spans within this gender group was carried out using a t-test. The relationships between body height and arm span were determined using simple correlation coefficients and their 95% confidence interval. Then linear regression analyses was performed to examine the extent to which arm span can reliably predict body height. Finally these relationships were plotted as scatter diagrams and regression lines. Statistical significance was set at  $p < 0.05$ .

## Results

A summary of the anthropometric measurements is shown in Table 1. The mean of the arm span for male subjects was  $182.53 \pm 7.53$  centimeters, which was  $2.03 \pm 1.50$  centimeters more than the body height and statistically insignificant ( $t = -1.604, p < 0.111$ ).

**Table 1.** Anthropometric Measurements of the Population

Body Height Range (Mean±SD)	Arm span Range (Mean±SD)
163.6-199.0 (182.53±7.53)	166.3-210.0 (184.55±9.03)

The simple correlation coefficient and their 95% confidence interval analysis between the anthropometric measurements are

presented in Table 2. The relationships between body height and arm span was high and significant in the sample.

**Table 2.** Correlation Between Body Height and Arm Span of the Study Subjects

Correlation Coefficient	95% confidence interval	Significance p-value
0.821	0.583–0.788	<0.000

The results of the linear regression analysis are shown in Table 3. The first of all models were derived by including age as a covariate. However, it was found that the contribution of age was insignificant and therefore the age was dropped and es-

timates were derived as univariate analysis. The high values of the regression coefficient signify that arm span significantly predicts body height in the applied sample.

**Table 3.** Results of Linear Regression Analysis Where the Arm Span Predicts the Body Height

Regression Coefficient	Standard Error (SE)	R-square (%)	t-value	p-value
0.821	4.323	67.5	13.272	0.000

The relationships between arm span measurements and body height among the above subjects is plotted as a scatter diagram.


## Discussion

The results of this study proved that Southern-Macedonian males are very tall with an average of 182.53 centimeters, but not taller than general male population in Montenegro with 183.21 centimeters (Quanjer et al., 2014) and Central- Monte-

negrins (Vujović, Bubanja, Tanase, & Milašinović, 2015). Parallel to this fact, they are still one of the tallest in Europe. It does not come close to 184.6 centimeters documented by Pineau et al. (Popović, Bjelica, Petković, Muratović, & Georgiev, 2014) and 183.8 centimeters of the Dutch male population measured in the last TNO nationwide survey in 2010 (cited in Popović, Bjelica, Tanase, & Milasinovic, 2015). From the results this study reached, probably, the tallest people might live on the northern part of Montenegro, mostly due to the reason the average body height in Central (Vujović et al., 2015) and Southern (this study) part of Montenegro did not reach the average body

height in whole Montenegro. On the other hand, the 182.53 centimeters average height of Southern-Montenegrin men is taller than 181.3 centimeters of the Lithuanians (Popović, Bjelica, & Hadžić, 2014), 180.9 centimeters of the Serbs (Grozdanov, 2011), 180.6 centimeters of the Icelanders (Bjelica et al., 2012) et many other nations. However, there is a hypothesis that Montenegrin males did not reach their full genetic potential yet, since they have been influenced by various environmental

factors (wars, poor economic situation, etc.) in the last few decades (Popović, Bjelica, Molnar, Jakšić, & Akpinar, 2013), while these influence was the smallest on the South part of Montenegro. Therefore, the authors believe that these circumstances had a negative bearing on the secular trend in Montenegro, while it is expected that the secular changes affecting height will go up in the following 20 years, comparing it to developed countries where this trend has already stopped.



**Figure 1.** Scatter Diagram and Relationship Between Arm Span Measurements and Body Height Among the Appled Sample

On the other hand, expectably, the arm span reliably predicts body height in this gender. However, the estimation equations which have been obtained in Southern-Montenegrins are different alike in general population, since arm span was closer

to body heights ( $2.03 \pm 1.50$  centimetres), more than in general population (Bjelica et al., 2012). This confirms the necessity for developing separate height models for each region in Montenegro.

## REFERENCES

- Bjelica, D., Popović, S., Kezunović, M., Petković, J., Jurak, G., & Grasgruber, P. (2012). Body Height and Its Estimation Utilizing Arm Span Measurements in Montenegrin Adults. *Anthropological Notebooks*, 18(2), 69–83.
- Coon, C. (1975). *The races of Europe*. Westport, Conn: Greenwood Press.
- Marfell-Jones, M., Olds, T., Stewart, A., & Carter, L. (2006). *International standards for anthropometric assessment*. Potchesfroom: International Society for the Advancement of Kinanthropometry.
- Quanjer, P. H., Capderou, A., Mazocioglu, M. M., Aggarwal, A., Popovic, S., Datta Banik, S., Tayie, F. A. K., Golshan, M., Ip, M. S. M., & Zelter, M. (2014). All-age relationship between arm span and height in different ethnic groups. *European Respiratory Journal*, 44, 905-912.
- Pineau, J. C., Delamarque, P., & Božinović, S. (2005). Average height of adolescents in the Dinaric Alps (in French). *Comptes Rendus Biologies* 328(9), 841–846.
- Popović, S., Bjelica, D., & Hadžić, R. (2014). Average body height of adolescents in Montenegro. In *Proceedings book of the 13th International Sport Sciences Congress (462-463)*. Konya: Selcuk University.
- Popović, S., Bjelica, D., Molnar, S., Jakšić, D., & Akpinar, S. (2013). Body Height and Its Estimation Utilizing Arm Span Measurements in Serbian Adults. *International Journal of Morphology*, 31(1), 271-279.
- Popović, S., Bjelica, D., Petković, J., Muratović, A., & Georgiev, G. (2014). Body Height and Its Estimation Utilizing Arm Span Measurements in Macedonian Adolescents. In *Abstract Book of the 7th Conference for Youth Sport (40)*, Ljubljana: Faculty of Sport, University of Ljubljana.
- Popović, S., Bjelica, D., Tanase, G. D., & Milasinovic, R. (2015). Body Height and Its Estimation Utilizing Arm Span Measurements in Bosnian and Herzegovinian Adults. *Montenegrin Journal of Sports Science and Medicine*, 4(1), 29-36.
- Vujović, D., Bubanja, M., Tanase, G. D., & Milašinović, R. (2015). Body Height and Its Estimation Utilizing Arm Span Measurements in Male Adolescents from Central Region in Montenegro. *Sport Mont*, 43-45/XII, 283-288.

R. Milašinović

University of Montenegro, Faculty for Sport and Physical Education, Narodne omladine bb, 81400 Nikšić, Montenegro  
e-mail: rajko.m@ac.me

